



Growth parameter estimates of *Listeria monocytogenes* in cooked chicken: effect of preparation of inoculum

Birk, Tina; Smith Ottosen, Sussi; Hansen, Tina Beck

Publication date:
2017

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Birk, T., Smith Ottosen, S., & Hansen, T. B. (2017). *Growth parameter estimates of Listeria monocytogenes in cooked chicken: effect of preparation of inoculum*. Poster session presented at 10th International Conference on Predictive Modelling in Food, Cordoba, Spain.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal




If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Growth parameter estimates of *Listeria monocytogenes* in cooked chicken: effect of preparation of inoculum

Tina Birk, Sussi Smith Ottosen and Tina Beck Hansen^a

Objective


Growth curves, and estimates of lag times and growth rates, for *Listeria monocytogenes* on slices of sous-vide cooked chicken breast were compared for four inoculum preparation procedures; coldBHI (control procedure), direct30 and direct8 (t-test against the control), and coldMRD (graphical comparison of lag times, λ , and growth rates, μ_{\max} , with the control). Inoculum volumes of 0.44 ± 0.11 % were used.



Conclusion




Even with inoculation volumes lower than 1%, fitted growth parameter estimates were affected by the preparation of inoculum.

- λ was mostly affected by growth phase then temperature shifts and lastly presence of nutrients, such as glucose, in the inoculum
- μ_{\max} was only affected by presence of nutrients



Results: cross-validation

Procedure	μ_{\max} model	Bias factor	Accuracy factor
coldBHI	$\sqrt{\mu_{\max}} = 0.0350 \cdot (T + 0.37)$	1.12	1.14
coldMRD	$\sqrt{\mu_{\max}} = 0.0344 \cdot (T - 0.21)$	0.88	1.17



a: for further information contact Tina Beck Hansen at tibha@food.dtu.dk, National Food Institute, Technical University of Denmark, Kemitorvet, Building 202, DK-2800 Kgs. Lyngby, Denmark